SEMICLASSICAL RESOLVENT BOUNDS FOR WEAKLY DECAYING POTENTIALS

JEFFREY GALKOWSKI AND JACOB SHAPIRO

Abstract. In this note, we prove weighted resolvent estimates for the semiclassical Schrödinger operator $h^2 + V(x) : L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$, $n \geq 2$. The potential V is real-valued, and assumed to either decay at infinity or to obey a radial α-Hölder continuity condition, $0 < \alpha \leq 1$, with sufficient decay of the local radial C^α norm toward infinity. Note, however, that in the Hölder case, the potential need not decay. If the dimension $n \geq 3$, the resolvent bound is of the form $\exp\left(Ch^{-\frac{1}{3+\alpha}}[(1-\alpha)\log(h^{-1}) + c]\right)$, while for $n = 1$ it is of the form $\exp(Ch^{-1})$. A new type of weight and phase function construction allows us to reduce the necessary decay even in the pure L^1 case.

1. Introduction and statement of results

Let $\Delta := \sum_{j=1}^n \partial_j^2 \geq 0$ be the Laplacian on $\mathbb{R}^n$, $n \geq 2$. In this article, we study the semiclassical Schrödinger operator

$$P(h) := h^2\Delta + V : L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n), \qquad h > 0,$$

where $V \in L^1_{\mathrm{loc}}(\mathbb{R}^n;\mathbb{R})$. We assume either that V satisfies a radial α-Hölder continuity condition, $0 < \alpha \leq 1$, or that it is only L^1 but decaying. When $n \geq 3$, we use $(r,\omega) = (|x|, x/|x|) \in (0,\infty) \times S^{n-1}$ to denote polar coordinates on $\mathbb{R}^n \setminus \{0\}$.

When V is only L^1, we assume

$$|V| \leq c_1\langle r\rangle^{-2}m(r), \tag{1.1}$$

for some

$$c_1 > 0, \qquad 0 < m(r) \leq 1, \qquad m(r)\langle r\rangle^{1/2} \in L^2(0,\infty), \tag{1.2}$$

and where $\langle x\rangle = \langle r\rangle := (1+r^2)^{1/2}$.

Since $V \in L^1_{\mathrm{loc}}(\mathbb{R}^n;\mathbb{R})$, by the Kato-Rellich Theorem, $P(h)$ is self-adjoint $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ with respect to the domain $H^2(\mathbb{R}^n)$. Therefore, the resolvent $(P - z)^{-1}$ is bounded $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ for all $z \in \mathbb{C} \setminus \mathbb{R}$, and we obtain

Theorem 1. Let $n \geq 3$, m as in (1.2), $c_1 > 0$ and $E > 0$. Then there are $C > 0$ and $h_0 \in (0,1]$ so that for all $s > 1/2$, there is $C_s > 0$ such that for all $V \in L^1_{\mathrm{loc}}(\mathbb{R}^n;\mathbb{R})$ satisfying (1.1),

$$g_s(h,\varepsilon) \leq C_s \exp\left(h^{-\frac{4}{3}}(C\log h^{-1} + C_s)\right), \qquad \varepsilon > 0,\ h \in (0,h_0], \tag{1.3}$$

where

$$g_s(h,\varepsilon) := \|\langle x\rangle^{-s}(P(h) - E - i\varepsilon)^{-1}\langle x\rangle^{-s}\|_{L^2(\mathbb{R}^n)\to L^2(\mathbb{R}^n)}. \tag{1.4}$$

This material is based upon work supported by the National Science Foundation under Grant No. 1440140, while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Fall semester of 2019. J. Shapiro was also supported in part by the Australian Research Council through grant DP180100589.

Now, we will assume throughout that $w^0;\ '^0 > 0$. Therefore, putting

$$:= \frac{'^{00}}{'^0} = (\log '^0)^0; \qquad W := \frac{w}{w^0} = \frac{1}{(\log w)^0}; \tag{2.9}$$

we calculate

$$\begin{aligned} A(r) \quad \frac{K}{2}B(r) &= w^0(E + ('^0)^2 \quad V_h) + w(2'^0'^{00} \quad V_h^0) \quad \frac{K}{2}\frac{(w(h^{\ 1}jR_hj + '^{00}))^2}{w^0 + 4h^{\ 1}'^0 w} \\ &= w^0\Big[E + ('^0)^2 \quad V_h + W(2'^0'^{00} \quad V_h^0) \quad \frac{K}{2}\frac{(w(h^{\ 1}jR_hj + '^{00}))^2}{w^{02} + 4h^{\ 1}'^0 ww^0}\Big] \\ &= w^0\Big[E + ('^0)^2(1 + 2W \ ) \quad V_h \quad WV_h^0 \quad \frac{K}{2}W^2\frac{((h^{\ 1}jR_hj + '^{00}))^2}{1 + 4h^{\ 1}'^0 W}\Big] \\ &\quad w^0\Big[E + ('^0)^2(1 + 2W \ ) \quad V_h \quad WV_h^0 \quad KW^2\frac{h^{\ 2}jR_hj^2 + ('^{00})^2}{1 + 4h^{\ 1}'^0 W}\Big]: \end{aligned}$$

Finally,

$$\begin{aligned} A(r) \quad \frac{K}{2}B(r) \quad w^0\Big[E + ('^0)^2(1 + 2W \quad KW^{\ 2}\min(W; \tfrac{h}{4'^0})) \\ V_h \quad WV_h^0 \quad KWh^{\ 2}jR_hj^2\min(W; \tfrac{h}{4'^0})\Big]: \end{aligned} \tag{2.10}$$

The key improvement in this article is that, to prove the main estimates (3.5) and (4.4), we work with W and rather than directly with w and $'$

Proof of Theorems 1, 2 and 3. Since increasing s in (1.4) decreases the resolvent norm, to prove (1.3), (1.9) and (1.13), we may assume without loss of generality that $0 < 2s \quad 1 < 1$.

Fix $= 2s \quad 1$. When $n \quad 3$, let $= \quad$ be as in (3.3). Let $'$, w, and $h_0 \ 2 \ (0;1]$ be

We then achieve (6.5) by replacing v by $\tilde{v}_k$ in (6.3) and sending $k \to 1$.

References

[Bu98] N. Burq. Decroissance de l'energie locale de l'equation des ondes pour le probleme exterieur et absence de resonance au voisinage du reel. *Acta Math.*, 180(1) (1998), 1–29

[Bu02] N. Burq. Lower bounds for shape resonances widths of long range Schrödinger operators. *Amer. J. Math.*, 124(4) (2002), 677–735

[CaVo02] F. Cardoso and G. Vodev. Uniform estimates of the resolvent of the Laplace-Beltrami operator on infinite volume Riemannian manifolds II. *Ann. Henri Poincare*, 3(4) (2002), 673–691

[DDZ15] K. Datchev, S. Dyatlov and M. Zworski. Resonances and lower resolvent bounds. *J. Spectr. Theory*, 5(3) (2015), 599–615

[Da14] K. Datchev. Quantitative limiting absorption principle in the semiclassical limit. *Geom. Funct. Anal.*, 24(3) (2014), 740–747

[DaSh19] K. Datchev and J. Shapiro. Semiclassical estimates for scattering on the real line, to appear in *Comm. Math. Phys.* arXiv 1903.02743

[KlVo19] F. Klopp and M. Vogel. Semiclassical resolvent estimate for bounded potentials. *Pure Appl. Anal.* (1)1 (2019), 1–25

[Sh19] J. Shapiro. Semiclassical resolvent bounds in dimension two. *Proc. Amer. Math. Soc.* 147(5) (2019), 1999–2008

[Sh17] J. Shapiro. Semiclassical resolvent bound for compactly supported L^1 potentials, to appear in *J. Spectr. Theory.* arXiv 1802.09008

[Vo14] G. Vodev. Semi-classical resolvent estimates and regions free of resonances. *Math. Nachr.* 287(7) (2014), 825–835

[Vo19a] G. Vodev. Semiclassical resolvent estimates for short-range L^1 potentials. *Pure Appl. Anal.* 1(2) (2019), 207–214

[Vo19b] G. Vodev. Semiclassical resolvent estimates for short-range L^1 potentials. II, to appear in *Asymptot. Anal.* arXiv:1901.01004

[Vo20a] G. Vodev. Semiclassical resolvents estimates for L^1 potentials on Riemannian manifolds. *Ann. Henri Poincare.* 21(2), 437–459

[Vo20b] G. Vodev. Semiclassical resolvent estimates for Hölder potentials, arXiv:2002.12853.

Department of Mathematics, University College London, London, UK
Email address: j.galkowski@ucl.ac.uk

Mathematical Sciences Institute, Australian National University, Acton, ACT, Australia
Email address: Jacob.Shapiro@anu.edu.au